

Clindamycin Phosphate and Benzoyl Peroxide Gel 1.2%/2.5%

PK-159-0
184



HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use **CLINDAMYCIN PHOSPHATE and BENZOYL PEROXIDE** safely and effectively. See all prescribing information for **CLINDAMYCIN PHOSPHATE and BENZOYL PEROXIDE**.

CLINDAMYCIN PHOSPHATE and BENZOYL PEROXIDE gel

CLINDAMYCIN PHOSPHATE and BENZOYL PEROXIDE gel is indicated for the treatment of acne vulgaris in patients 12 years of age or older.

INDICATIONS AND USAGE

CLINDAMYCIN PHOSPHATE and BENZOYL PEROXIDE gel is indicated for the treatment of acne vulgaris in patients 12 years of age or older.

CONTRAINDICATIONS

Do not use **CLINDAMYCIN PHOSPHATE and BENZOYL PEROXIDE gel** if you are allergic to clindamycin, benzoyl peroxide, or any of the other ingredients.

WARNINGS AND PRECAUTIONS

Do not use **CLINDAMYCIN PHOSPHATE and BENZOYL PEROXIDE gel** if you are allergic to clindamycin, benzoyl peroxide, or any of the other ingredients.

ADVERSE REACTIONS

The following adverse reactions are common in patients 12 years of age or older who use **CLINDAMYCIN PHOSPHATE and BENZOYL PEROXIDE gel**.

DRUG INTERACTIONS

Do not use **CLINDAMYCIN PHOSPHATE and BENZOYL PEROXIDE gel** if you are allergic to clindamycin, benzoyl peroxide, or any of the other ingredients.

HOW TO USE CLINDAMYCIN PHOSPHATE and BENZOYL PEROXIDE gel

Apply a pea-sized amount of **CLINDAMYCIN PHOSPHATE and BENZOYL PEROXIDE gel** to the affected area of your face twice daily.

HOW TO STORE CLINDAMYCIN PHOSPHATE and BENZOYL PEROXIDE gel

Store **CLINDAMYCIN PHOSPHATE and BENZOYL PEROXIDE gel** at room temperature (20° to 25°C).

CONTENTS

Each box contains 30 tubes of **CLINDAMYCIN PHOSPHATE and BENZOYL PEROXIDE gel**.

PATIENT INFORMATION

CLINDAMYCIN PHOSPHATE and BENZOYL PEROXIDE gel

IMPORTANT: For use on skin only (topical use). Do not get Clindamycin Phosphate and Benzoyl Peroxide Gel in your mouth, eyes, or vagina or on your lips.

Read the Patient Information that comes with clindamycin phosphate and benzoyl peroxide gel each time you use it. There may be new information. This information does not take the place of talking with your doctor about your medical condition or your treatment.

What is clindamycin phosphate and benzoyl peroxide gel?

Clindamycin phosphate and benzoyl peroxide gel is a prescription medicine used to treat acne vulgaris. Clindamycin phosphate and benzoyl peroxide gel contains clindamycin phosphate and benzoyl peroxide. It is not known if clindamycin phosphate and benzoyl peroxide gel is safe and effective for use longer than 12 weeks. It is not known if clindamycin phosphate and benzoyl peroxide gel is safe and effective in children under 12 years of age.

Who should not use clindamycin phosphate and benzoyl peroxide gel?

Do not use clindamycin phosphate and benzoyl peroxide gel if you have:

- been allergic to clindamycin, benzoyl peroxide, or any of the other ingredients.
- had inflammation of the colon (colitis), or severe diarrhea with past antibiotic use.
- had your doctor tell you not to use it.

What should I tell my doctor before using clindamycin phosphate and benzoyl peroxide gel?

Tell your doctor about all the medicines and skin products you use. Especially tell your doctor if you will have surgery with general anesthesia. One of the medicines in clindamycin phosphate and benzoyl peroxide gel (clindamycin) can affect how certain medicines work when used together.

How to use clindamycin phosphate and benzoyl peroxide gel

Apply a pea-sized amount of clindamycin phosphate and benzoyl peroxide gel to the affected area of your face twice daily. Do not use more than the amount of clindamycin phosphate and benzoyl peroxide gel that your doctor tells you to use.

How to store clindamycin phosphate and benzoyl peroxide gel

Store clindamycin phosphate and benzoyl peroxide gel at room temperature (20° to 25°C). Do not use clindamycin phosphate and benzoyl peroxide gel if the container is open for more than 3 months.

Other information

Keep this and all other medicines out of the reach of children. This medicine should be kept out of the reach of children. This medicine should be kept out of the reach of children.

CLINDAMYCIN PHOSPHATE and BENZOYL PEROXIDE GEL

CLINDAMYCIN PHOSPHATE and BENZOYL PEROXIDE GEL

IMPORTANT: For use on skin only (topical use). Do not get Clindamycin Phosphate and Benzoyl Peroxide Gel in your mouth, eyes, or vagina or on your lips.

Read the Patient Information that comes with clindamycin phosphate and benzoyl peroxide gel each time you use it. There may be new information. This information does not take the place of talking with your doctor about your medical condition or your treatment.

What is clindamycin phosphate and benzoyl peroxide gel?

Clindamycin phosphate and benzoyl peroxide gel is a prescription medicine used to treat acne vulgaris. Clindamycin phosphate and benzoyl peroxide gel contains clindamycin phosphate and benzoyl peroxide. It is not known if clindamycin phosphate and benzoyl peroxide gel is safe and effective for use longer than 12 weeks. It is not known if clindamycin phosphate and benzoyl peroxide gel is safe and effective in children under 12 years of age.

Who should not use clindamycin phosphate and benzoyl peroxide gel?

Do not use clindamycin phosphate and benzoyl peroxide gel if you have:

- been allergic to clindamycin, benzoyl peroxide, or any of the other ingredients.
- had inflammation of the colon (colitis), or severe diarrhea with past antibiotic use.
- had your doctor tell you not to use it.

What should I tell my doctor before using clindamycin phosphate and benzoyl peroxide gel?

Tell your doctor about all the medicines and skin products you use. Especially tell your doctor if you will have surgery with general anesthesia. One of the medicines in clindamycin phosphate and benzoyl peroxide gel (clindamycin) can affect how certain medicines work when used together.

How to use clindamycin phosphate and benzoyl peroxide gel

Apply a pea-sized amount of clindamycin phosphate and benzoyl peroxide gel to the affected area of your face twice daily. Do not use more than the amount of clindamycin phosphate and benzoyl peroxide gel that your doctor tells you to use.

How to store clindamycin phosphate and benzoyl peroxide gel

Store clindamycin phosphate and benzoyl peroxide gel at room temperature (20° to 25°C). Do not use clindamycin phosphate and benzoyl peroxide gel if the container is open for more than 3 months.

Other information

Keep this and all other medicines out of the reach of children. This medicine should be kept out of the reach of children. This medicine should be kept out of the reach of children.

Clindamycin Phosphate and Benzoyl Peroxide Gel	Benzoyl Peroxide		Clindamycin Phosphate		Benzoyl Peroxide and Clindamycin Phosphate	
	1.2%	2.5%	1.2%	2.5%	1.2%	2.5%
Common	22%	22%	10%	10%	10%	10%
Uncommon	6%	6%	1%	1%	1%	1%
Rare	1%	1%	0%	0%	0%	0%
Very Rare	0%	0%	0%	0%	0%	0%
Not Known	0%	0%	0%	0%	0%	0%



